



A Union of Professionals

COVID-19 Resources:

Here is **evidence to help you make your case** for respiratory protection for members caring for suspected and confirmed COVID-19 patients

APRIL 16, 2020

Use this resource if the employer claims that COVID-19 is primarily spread through droplet and contact transmission and is not aerosolized.

When the Centers for Disease Control and Prevention changed its infection control guidance, recommending facemasks instead of N95 respirators for those providing care to suspected and confirmed COVID-19 patients, the agency justified the action based on concern about the supply and supply chains for N95 production. But the **CDC also claimed without evidence that SARS-CoV-2 is also primarily spread via droplet and contact transmission.** In seeming contradiction, the agency still recommends that workers use N95 respirators when the supply chain issues are resolved.

Hospitals and other healthcare employers have been quick to use the argument that respiratory protection is not needed rather than finding other sources of respiratory protection, including other kinds of filtering face piece respirators (N99s, N100s, P100s, elastomers) and powered air-purifying respirators (PAPRs). Many employers have immediately moved from conventional capacity for providing healthcare workers with

respiratory protection to crisis capacity, denying it completely, putting healthcare workers' health and ability to provide care in jeopardy.

Much has to be done to increase the production and supply of N95 respirators, including enactment of the Defense Production Act to promote immediate manufacture of respirators. Employers need to invest in reusable respirators.

If the employer is claiming that respirators are not needed as per the CDC, there is growing evidence that the virus that causes COVID-19 is aerosolized over short distances and persists for long periods of time. The traditional understanding of infectious disease transmission (contact, droplet and airborne) is 70 years old. There is a growing body of research indicating that many viruses, including coronaviruses, are aerosolized when an infected person speaks and exhales. Surgical masks only protect the wearer from droplets caused by coughs and sneezes.

Find these resources and more at www.aft.org/coronavirus

The **American Federation of Teachers** is a union of 1.7 million professionals that champions fairness; democracy; economic opportunity; and high-quality public education, healthcare and public services for our students, their families and our communities. We are committed to advancing these principles through community engagement, organizing, collective bargaining and political activism, and especially through the work our members do.

Randi Weingarten
PRESIDENT

Loretta Johnson
SECRETARY-TREASURER

Evelyn DeJesus
EXECUTIVE VICE PRESIDENT



A good explanation is found here:

"COVID-19 transmission messages should hinge on science"

Brosseau, L. March 16, 2020

bit.ly/covid19brosseau

Here is a sampling of the research that has been done since COVID-19 emerged. Many of the studies are small and have not been peer-reviewed yet, but they were published in reputable journals in order to share knowledge while people need the information.

1. In a study from the National Institutes of Health involving scientists from CDC, UCLA and Princeton, researchers found that SARS-CoV-2 virus was detectable in the air for up to three hours.

"Aerosol and Surface Stability of SARS-CoV-2 as Compared with "SARS-CoV-1," *New England Journal of Medicine*, Doremalen, N. et al., March 17, 2020.

bit.ly/aerosol-cv19

2. During the initial isolation of 13 individuals confirmed positive with COVID-19 infection, air and surface samples were collected in 11 isolation rooms to examine viral shedding from isolated individuals. SARS-CoV-2 is shed during respiration, toileting, and fomite contact, indicating that infection may occur in both direct and indirect contact, supporting the need for airborne isolation precautions.

"Transmission Potential of SARS-CoV-2 in Viral Shedding Observed at the University of Nebraska Medical Center," Santarpia, J. et al., preprint publication, March 23, 2020

bit.ly/covid19santarpia

3. In a small German study, researchers found that the novel coronavirus quickly begins producing high viral loads, sheds efficiently, and grows well in the nose, mouth, nasal cavity and throat.

"Virological assessment of hospitalized cases of coronavirus disease 2019," Wölfel, R. et al., *Annals of Internal Medicine*, March 5, 2020.

bit.ly/covid19wolfel

4. Researchers in Wuhan, China, measured the concentration and aerodynamic characteristics of airborne SARS-CoV-2 aerosol in different areas of two hospitals. The findings add support to the hypothesis that virus-laden aerosol deposition may play a role in surface contamination and subsequent contact transmission, particularly on toilets and in areas without negative air pressure and/or frequent air exchanges.

"Aerodynamic Characteristics and RNA Concentration of SARS-CoV-2 Aerosol in Wuhan Hospitals during COVID-19 Outbreak," Liu, Y. et al., March 8, 2020.

bit.ly/covid19liu

5. When oxygen is delivered through nasal catheter, mask or non-invasive ventilation, substantial exhaled air is released into the air, which can increase dispersion of the virus, and subsequently increase the risk of nosocomial infection.

"More awareness is needed for severe acute respiratory syndrome coronavirus 2019 transmission through exhaled air during non-invasive respiratory support: experience from China," Guan, L. et al. *European Respiratory Journal*, Feb. 18, 2020.

bit.ly/covid19guan

6. Wuhan researchers found N95 use protected healthcare workers from infection compared to healthcare workers who did not use N95s. Includes air sampling data.

Wang, X., Pan, Z., & Cheng, Z. (2020). "Association Between 2019-nCoV Transmission and N95 Respirator Use" *medRxiv*.

bit.ly/covid19wang

7. "Recognition of Aerosol Transmission of Infectious Agents: a Commentary"

This scholarly article explains the distinctions among droplet, airborne and aerosol transmission. Tellier et al. *BMC Infectious Diseases*, 2019.

(bit.ly/covid19tellier)

8. "Unmasked: Experts explain necessary respiratory protection for COVID-19"

(bit.ly/CIDRAP-respiratoryprotection)

9. Researchers from Wuhan collected indoor air samples in a COVID-19 intensive care unit and a general ward in a Wuhan hospital, as well as collecting swab samples. The virus was detected approximately four meters from patients and was widely distributed on high touch surfaces, floors and caregivers' shoes.

"Aerosol and Surface Distribution of Severe Acute Respiratory Syndrome Coronavirus 2 in Hospital Wards, Wuhan, China, 2020," Zhen-Dong Guo et al. *Emerging Infectious Diseases*, Centers for Disease Control, early release July 2020

(bit.ly/covid19-surfaces)